Author Index

Abban E. K., *see* Oberst O., 235.
Abellan E., *see* Nathanailides C., 885.
Adler P. R., *see* Heinen J. M., 699.
Agnese J.-F., *see* Otémé Z. J., 805.
Agustin L. Q., *see* Macaranas J. M., 597.
Al Hinty H. M., *see* Siddiqui A. Q., 515.
Al-Amoudi M., *see* El-Sayed A.-F. M., 681.
Al-Asgah N. A., *see* Al-Ogaily S. M., 523.
Al-Ogaily S. M., Al-Asgah N. A. & Ali A.: Effect of feeding different grain sources on the growth performance and body composition of tilapia, *Oreochromis niloticus* (L.), 523.
Al-Owafeir M. A. & Belal I. E. H.: Replacing alm oil for soybean oil in tilapia, *Oreochromis niloticus* (L.), feed, 221.
Alam M. J., Ang K. J. & Begum M.: Ingestion efficiency of *Macrobranchium rosenbergii* (de Man) larvae feeding on Artemia, *Monia micrura* Kurtz and their combination, 113.
Alam M. K., Maughan O. E. & Matter W. J.: Growth response of indigenous and exotic carp species to different protein sources in pelleted feeds, 673.
Alanärä A. & Kiessling A.: Changes in demand feeding behaviour in Arctic charr, *Salvelinus alpinus* L., caused by differences in dietary energy content and reward level, 479.
Alexis M. N., *see* Nengas I., 147.
Ali A., *see* Al-Ogaily S. M., 523.
Ali S. A., *see* Siddiqui A. Q., 515.
Ambak M. A., *see* Khan M. S., 823.
Ang K. J., *see* Alam M. J., 113; *see also* Das N. N., 543; *see also* Khan M. S., 823.
Audet C., *see* Dumas S., 245.
Awaïss A., Kestemont P. & Micha J. C.: Fatty acid profiles of two freshwater fish larvae (gudgeon and perch) reared with *Brachionus calyciflorus* Pallas (rotifer) and/or dry diet, 651.

Baardvik B. M., *see* Siikavuopio S. I., 205.
Baker R. T. M. & Davies S. J.: Oxidative nutritional stress associated with feeding rancid oils to African catfish, *Clarias gariepinus* (Burchell) and the protective role of α-tocopherol, 795.
Balebona M. C., *see* Sedano J., 323.
Basu T. K. & Bhattacharyya B.: Application of Hill's procedure for growth curve analysis of a freshwater major carp of India, *Catla catla* (Ham), during their early developmental period, 891.
Battaglene S. C. & Selosse P. M.: Hormone-induced ovulation and spawning of captive and wild broodfish of the catadromous Australian bass, *Macquaria novemaculeata* (Steindachner), (Percichthyidae), 191.
Begum M., *see* Alam M. J., 113.
Belal I. E. H., *see* Al-Owafeir M. A., 221.
Beveridge M. C. M., *see* Chellappa S., 91; *see also* Chellappa S., 945.
Bhatnagar A., *see* Garg S. K., 157.
Bhattacharyya B., *see* Basu T. K., 891.
Blanc J. M., *see* Dumas S., 245.
Boon J. H., *see* Hariati A. M., 255; *see also* Hariati A. M., 277; *see also* Peutz I. L. J. A., 437.
Bordas M. A., *see* Sedano J., 323.
Borrego J. J., *see* Sedano J., 323.
Briggs M. R. P. & Funge-Smith S. J.: The effects of zeolites and other alumino-silicate clays on water quality at various salinites, 301.
Briggs M. R. P. & Funge-Smith S. J.: The potential use of *Gracilaria* sp. meal in diets for juvenile *Penaeus monodon* Fabricius, 345.
Brown S. E. & Goulder R.: Extracellular-enzyme activity in trout-farm effluents and a recipient river, 895.
Brummett R. E. & Katambalika K.: Protocols for the development of indigenous species: polyculture of indigenous species under Malawian smallholder conditions, 225.
Brunsvik P. S., Finstad B. & Nilssen N. J.: Influence of sea ranching on hatcdhery-reared anadromous Arctic charr, *Salvelinus alpinus* (L.), 735.
Butler A. & Cross T. F.: Genetic differences between successive year classes of two strains of reared rainbow trout, *Oncorhynchus mykiss* (Walbaum), 643.
Buttle L. G., Uglow R. F. & Cowx I. G.: Temperature and dietary factors affecting the nitrogen efflux rates of the African catfish, *Clarias gariepinus* (Burchell), 391.
Bye J. A. T., *see* Strutton P. G., 497.

Cacho M. S. R. F., *see* Chellappa S., 91.
Carmona-Osalde C., *see* Rodríguez-Serna M., 67.
Chandran M. R., *see* Sivakami R., 375.
Chavez-Sanchez M. C., *see* Martinez-Palacios C. A., 455.
Chellappa N. T., *see* Chellappa S., 945.
Chellappa S., Cacho M. S. R. F., Huntingford F. A. & Beveridge M. C. M.: Observations on induced breeding of the Amazonian fish tambaqui, *Colossoma macropomum* (Cuvier), using CPE and HCG treatments, 91.
Chellappa S., Chellappa N. T., Silva E. A., Huntingford F. A. & Beveridge M. C. M.: The diet of hybrid red tilapia, *Oreochromis niloticus* (L.) × *Oreochromis mossambicus* (Peters), reared in the freshwater ponds of north-eastern Brazil, 945.
Ciereszko A., *see* Dabrowski K., 539.
Cohen S., *see* Karplus I., 447.
Conides A. J., *see* Klaoudatos S. D., 765.
Costello M. J., *see* Papoutsoglou S., 25.
Cowx I. G., *see* Buttle L. G., 391.
Cross T. F., *see* Butler A., 643; *see also* Rogan E., 9.
Cui G. H., *see* Guan R. Z., 587.

Dabrowski K. & Ciereszko A.: The dynamics of gonad growth and ascorbate status in yellow perch, *Perca flavescens* (Mitchill), 539.

© 1996 Blackwell Science Ltd.

Dabrowski K., *see* Moreau R., 953.
Das N. N., Saad C. R., Ang K. J., Law A. T. & Harmin S. A.: Diet formulation of *Macrobranchium rosenbergii* (de Man) broodstock based on essential amino acid profile of its eggs, 543.
Davies S. J., *see* Baker R. T. M., 795; *see also* Henrique M. M. F., 405; *see also* Nengas I., 147; *see* Ojaveer H., 603;
Day O. J., Jones D. A. & Howell B. R.: Food consumption, growth and respiration of sole, *Solea solea* (L.), during early ontogeny in a hatchery environment, 831.
de la Noüe J., *see* Dumas S., 245.
del Valle G., Taniguchi N. & Tsujimura A.: Genetic differences in some haematological traits of communally reared clonal ayu, *Plecoglossus altivelis* Teminck & Schlegel, under stressed and non-stressed conditions, 787.
Demska-Zakes K., *see* Zakes Z., 841.
Dick T. A., *see* Wangila-B. C. C., 35.
Dumas S., Blanc J. M., Vallée F., Audet C. & de la Noüe J.: Survival, growth, sexual maturation and reproduction of brook charr, *Salvelinus fontinalis* (Mitchill), Arctic charr, *Salvelinus alpinus* L., and their hybrids, 245.

Edwards D. B., *see* Nugues M. M., 913; *see also* Spencer B. E., 261.
Ekanem S. B.: Effects of feeding frequency, moist and dry feeds on the growth of *Chrysichthys nigrodigitatus* Lacépède and on pond water quality, 107.
Eknath A. E., *see* Macaranas J. M., 597.
El-Ghobashy A., *see* El-Sayed A.-F. M., 681.
El-Sayed A.-F. M., El-Ghobashy A. & Al-Amoudi M.: Effects of pond depth and water temperature on the growth, mortality and body composition of Nile tilapia, *Oreochromis niloticus* (L.), 681.
English W. R., *see* Sureshwaran S., 557.

Fanning G. R., *see* Sahul Hameed A. S., 853.
Farmer J. J., *see* Sahul Hameed A. S., 853.
Felton S. P., Landolt M. L., Grace R. & Palmisano A.: Effects of selenium dietary enhancement on hatchery-reared coho salmon, *Oncorhynchus kisutch* (Walbaum), when compared with wild coho: hepatic enzymes and seawater adaptation evaluated, 135.
Feng X. H., *see* Guan R. Z., 587.
Finstad B., *see* Brunsvik P. S., 735.
Forsberg O. I.: Ammonia excretion rates from post-smolt Atlantic salmon, *Salmo salar* L., in land-based farms, 937.
Frazer A. W., *see* Heasman M. P., 505.
Funge-Smith S. J., *see* Briggs M. R. P., 301; *see also* Briggs M. R. P., 345.

Garcia-Abiado M. A. R., Merculio J. G. & Mair G. C.: The effect of lutenizing hormone-releasing hormone analogue (LHRHa) on sperm count of *Oreochromis niloticus* (L.), 95.
Garcia-Riera M. P. & Hemre G.-I.: Organ responses to ^{14}C-glucose injection in Atlantic halibut, *Hippoglossus hippoglossus* (L.), acclimated to diet of varying carbohydrate content, 565.
Garg S. K. & Bhatnagar A.: Effect of varying doses of organic and inorganic fertilizers on plankton production and fish biomass in brackish water fish ponds, 157.
Gasca-Leyva E., *see* Olvera-Novoa M. A., 167.
Gelman E., *see* Rothbard S., 175.
Gislason G., Olsen R. E. & Ringo E.: Comparative effects of dietary Na^+-lactate on Arctic charr, *Salvelinus alpinus* L., and Atlantic salmon, *Salmo salar* L.,429.
Gjerde B., *see* Rye M., 121.
Goard L. G., *see* Nell J. A., 689.
Gomez-Muñoz V. M., *see* Hernandez-Llamas A., 711.
Goodall E. A., *see* Menzies F. D., 183.
Goulder R., *see* Brown S. E., 895.
Grace R., *see* Felton S. P., 135.
Guan R. Z., Zhou L. H., Cui G. H. & Feng X. H.: Studies on the artificial propagation of *Monopterus albus* (Zuiew), 587.
Gulbrandsen J., Lein I. & Holmefjord I.: Effects of light administration and algae in first feeding of Atlantic halibut larvae, *Hippoglossus hippoglossus* (L.), 101.

Hand R. E., *see* Nell J. A., 689.
Hankins J. A., Heinen J. M., 699.
Hariati A. M., Wiadnya D. G. R., Sukkel M., Boon J. H. & Verdegem M. C. J.: Pond production of *Penaeus monodon* in relation to stocking density, survival rate and mena weight at harvest in East Java, Indonesia, 277.
Hariati A. M., Wiadnya D. G. R., Tanck M. W. T., Boon J. H. & Verdegem M. C. J.: *Penaeus monodon* (Fabricius) production related to water quality in East Java, Indonesia, 255.
Harmin S. A., *see* Das N. N., 543.
Harris S. A. & Probyn T.: Nitrogen extraction and absorption efficiencies of white steenbras, *Lithognathus lithognathus* Cuvier (sparidae), under experimental culture conditions, 43.
Heasman M. P., O'Connor W. A. & Frazer A. W.: Effects of fertilization and incubation factors on the quality and yield of scallop, *Pecten fumatus* Reeve, larvae, 505.
Hecht T., *see* Weyl O., 757.
Heggberget T. G., Økland F. & Ugedal O.: Prespawning migratory behaviour of wild and farmed Atlantic salmon, *Salmo salar* L., in a north Norwegian river, 313.
Heinen J. M., Hankins J. A. & Adler P. R.: Water quality and waste production in a recirculating trout-culture system with feeding of a higher-energy or a lower-energy diet, 699.
Hem S., *see* Otémé Z. J., 805.
Hemre G.-I., *see* Garcia-Riera M. P., 565.
Henricson J., *see* Näslund I., 815.
Henrique M. M. F., Morris P. C. & Davies S. J.: Vitamin C status and physiological response of the gilthead sea bream, *Sparus aurata* L., to stressors associated with aquaculture, 405.
Hernandez-Llamas A. & Gomez-Muñoz V. M.: Growth and survival response of the Catarina scallop, *Argopecten circularis* (Sowerby), to stocking density and length of culture period, 711.
Hickman-Brenner F. W., *see* Sahul Hameed A. S., 853.

© 1996 Blackwell Science Ltd.

Himelman J. H., *see* Lodeiros C. J. M., 749.
Holmefjord I., *see* Gulbrandsen J., 101.
Horney B. S., *see* Peutz I. L. J. A., 437.
Howell B. R., *see* Day O. J., 831.
Huisman E. A., *see* van Dam A. A., 463; *see also* van der Meer M. B., 531.
Huntingford F. A., *see* Chellappa S., 91; *see also* Chellappa S., 945.

Jana B. B., *see* Sahu S. N., 931.
Jobling M., *see* Siikavuopio S. I., 205.
Johnson G. R., *see* Peutz I. L. J. A., 437.
Jones D. A., *see* Day O. J., 831.

Kaiser H., *see* Weyl O., 757.
Kaiser M. J., *see* Nugues M. M., 913; *see also* Spencer B. E., 261.
Kamstra A., Span J. A. & van Weerd J. H.: The acute toxicity and sublethal effects of nitrite on growth and feed utilization of European eel, *Anguilla anguilla* (L.), 903.
Karplus I., Milstein A., Cohen S. & Ross L. G.: The effect of stocking different ratios of common carp, *Cyprinus carpio* L., and tilapias in polyculture ponds on production characteristics and profitability, 447.
Katambalika K., *see* Brummett R. E., 225.
Kathiresan K., *see* Rajendra N., 745.
Kestemont P., *see* Awaïss A., 651.
Khalil A. M.: The influence of algal concentration and body size on filtration and ingestion rates of the clam, *Tapes decussatus* (L.), (Mollusca: Bivalvia), 613.
Khan M. S., Ang K. J. & Ambak M. A.: The effect of varying dietary protein level on the growth, food conversion, protein utilization and body composition of tropical catfish, *Mystus nemurus* (Cuvier & Valenciennes), cultured in a static pond water system, 823.
Kiessling A., *see* Alanärä A., 479.
Klaoudatos S. D. & Conides A. J.: Growth, food conversion, maintenance and long-term survival of gilthead sea bream, *Sparus auratus* L., juveniles after abrupt transfer to low salinity, 765.
Knights B.: Studies of feeding stimulation in young eels, *Anguilla anguilla* L., before and after first-feeding using a novel rapid-screening method, 379.
Kouassi C. K., *see* Otémé Z. J., 805.
Kristjánsson L. Th. & Vøllestad L. A.: Individual variation in progeny size and quality in rainbow trout, *Oncorhynchus mykiss* (Walbaum), 335.
Kucharczyk D., Kujawa R., Mamcarz A., Skrzypczak A. & Wyszomirska E.: Induced spawning in perch, *Perca fluviatilis* L., using carp pituitary extract and HCG, 847.
Kujawa R., *see* Kucharczyk D., 847.

Lahnsteiner F., Patzner R. A. & Weismann T.: Semen cryopreservation of salmonid fishes: influence of handling parameters on the postthaw fertilization rate, 659.
Laing I., *see* Utting S. D., 573.
Lam T. J., *see* Sivaloganathan B., 635.
Landolt M. L., *see* Felton S. P., 135.
Law A. T., *see* Das N. N., 543.
Lein I., *see* Gulbrandsen J., 101.
Leung P. S., *see* Ling B. H., 775.
Ling B. H., Leung P. S. & Sang Y. C.: Export performance of major cultured shrimp producers in the Japanese and US markets, 775.
Lodeiros C. J. M. & Himelman J. H.: Influence of fouling on the growth and survival of the tropical scallop, *Euvola (Pecten) ziczac* (L., 1758) in suspended culture, 749.
Lopez-Albors O., *see* Nathanailides C., 885.
Loy G. L., *see* Sivaloganathan B., 635.

Macaranas J. M., Agustin L. Q. & Eknath A. E.: Multiple haemoglobins in three tilapiine species of the genus *Oreochromis* and in eight strains of *O. niloticus* (L.), 597.
Maguire G. B., *see* Nell J. A., 689.
Mair G. C., Garcia-Abiado M. A. R., 95.
Mamcarz A., *see* Kucharczyk D., 847.
Martinez-Palacios C. A., Chavez-Sanchez M. C. & Ross L. G.: The effects of water temperature on food intake, growth and body composition of *Cichlasoma urophthalmus* (Günther) juveniles, 455.
Martinez-Palacios C. A., Ross L. G. & Sanchez Licea V. H.: The tolerance to salinity, respiratory characteristics and potential for aquaculture of the Central American cichilid, *Cichlasoma synspilum* (Hubbs, 1935), 215.
Martinez-Palacios C. A., *see* Olvera-Novoa M. A., 167.
Matter W. J., *see* Alam M. K., 673.
Maughan O. E., *see* Alam M. K., 673.
McAdam S. P., *see* Nell J. A., 689.
McLoughlin M. F., *see* Menzies F. D., 183.
Menzies F. D., McLoughlin M. F., Wheatley S. B. & Goodall E. A.: Development of a computerized information retrieval system for Atlantic salmon, *Salmo salar* L., production, 183.
Merculio J. G., *see* Garcia-Abiado M. A. R., 95.
Micha J. C., *see* Awaïss A., 651.
Millican P. F., *see* Utting S. D., 573.
Milstein A. & Svirsky F.: Effect of fish species combinations on water chemistry and plankton composition in earthen fish ponds, 79.
Milstein A., *see* Karplus I., 447.
Mitchell J. G., *see* Strutton P. G., 497.
Moksness E. & Pavlov D. A.: Management by life cycle of wolffish, *Anarhichas lupus* L., a new species for coldwater aquaculture: a technical paper, 865.
Moksness E., *see* Pavlov D. A., 421; *see also* Steinarsson A., 925.
Moreau R. & Dabrowski K.: Feeding stimulants in semipurified diets for juvenile lake sturgeon, *Acipenser fulvescens* Rafinesque, 953.
Morgan R. I. G. & Paveley D. S.: A simple batch mark for fish studies using injected elastomer, 631.
Morinigo M. A., *see* Sedano J., 323.
Morris P. C., *see* Henrique M. M. F., 405; *see* Ojaveer H., 603.
Muir J. F., *see* Young J. A., 283.
Mukhopadhyay P. K. & Rout S. K.: Effects of different dietary lipids on growth and tissue fatty acid changes in fry of the carp, *Catla catla* (Hamilton), 623.

Näslund I. & Henricson J.: Growth and sexual maturation of six stocks of brown trout, *Salmo trutta* L., in culture, 815.
Nathanailides C., Lopez-Albors O., Abellan E., Vazquez J. M., Tyler D. D., Rowlerson A. & Strickland N. C.: Muscle cellularity in relation to somatic growth in the European sea bass, *Dicentrarchus labrax* (L.), 885.
Nell J. A., Hand R. E., Goard L. G., McAdam S. P. & Maguire G. B.: Studies of triploid oysters in Australia: evaluation of cytochalasin B and 6-dimethylamino-purine for triploidy induction in Sydney rock oysters, *Saccostrea commercialis* (Iredale & Roughley), 689.
Nengas I., Alexis M. N. & Davies S. J.: Partial substitution of fishmeal with soybean meal products and derivatives in diets for the gilthead sea bream, *Sparus aurata* (L.), 147.
Nilssen N. J., *see* Brunsvik P. S., 735.
Nugues M. M., Kaiser M. J., Spenser B. E. & Edwards D. B.: Benthic community changes associated with intertidal oyster cultivation, 913.
Nunez Rodriguez J., *see* Otémé Z. J., 805.

O'Connor W. A., *see* Heasman M. P., 505.
Oberst O., Abban E. K. & Vilwock W.: Biochemical and immunological markers for the discrimination of three *Tilapia* species: *T. zilli* Gervais, *T. guineensis* Bleeker and *T. dageti* Thys v.d. Audenaerde (Pisces: Cichlidae) from West Africa, 235.
Ojaveer H., Morris P. C., Davies S. J. & Russell P.: The response of thick-lipped grey mullet, *Chelon labrosus* (Risso), to diets of varied protein-to-energy ratio, 603.
Økland F., *see* Heggberget T. G., 313.
Olvera-Novoa M. A., *see* Rodríguez-Serna M., 67.
Olvera-Novoa M. A., Gasca-Leyva E. & Martinez-Palacios C. A.: The dietary protein requirements of *Cichlasoma synspilum* Hubbs, 1935 (Pisces: Cichlidae) fry, 167.
Olsen R. E., *see* Gislason G., 429.
Oorschot R. W. A., *see* Peutz I. L. J. A., 437.
Otémé Z. J., Nunez Rodriguez J., Kouassi C. K., Hem S. & Agnese J.-F.: Testicular structure, spermatogenesis and sperm cryopreservation in the African clariid catfish, *Heterobranchus longifilis* (Valenciennces, 1840), 805.

Palmisano A., *see* Felton S. P., 135.
Papoutsoglou S., Costello M. J., Stamou E. & Tziha G.: Environmental conditions at sea-cages, and ectoparasites on farmed European sea bass, *Dicentrarchus labrax* (L.), and gilthead sea bream, *Sparus aurata* L., at two farms in Greece, 25.
Patzner R. A., *see* Lahnsteiner F., 659.
Paveley D. S., *see* Morgan R. I. G., 631.
Pavlov D. A. & Moksness E.: Swelling of wolffish, *Anarhichas lupus* L., eggs and prevention of their adhesiveness, 421.
Pavlov D. A., *see* Moksness E., 865.
Petridis D. & Rogdakis I.: The development of growth and feeding equations for sea bream, *Sparus aurata* L., culture, 413.
Peutz I. L. J. A., Oorschot R. W. A., Johnson G. R., Horney B. S. & Boon J. H.: The leucogram as an indicator of marinecultured rainbow trout, *Oncorhynchus mykiss* (Walbaum), health in The Netherlands, 437.
Premkishore G., *see* Sivakami R., 375.
Probyn T., *see* Harris S. A., 43.
Puigcerver M.: Analysis on individual growth of juvenile carpet shell clams, *Tapes decussatus* (L.), in three different rearing conditions employed in the Ebro's Delta Spain, 399.

Rabbinge R., *see* van Dam A. A., 463.
Rajendra N. & Kathiresan K.: Effect of effluent from a shrimp pond on shoot biomass of mangrove seedlings, 745.
Rao P. V., *see* Sahul Hameed A. S., 853.
Ringo E., *see* Gislason G., 429.
Rodríguez-Serna M., Olvera-Novoa M. A. & Carmona-Osalde C.: Nutritional value of animal by-product meal in practical diets for Nile tilapia, *Oreochromis niloticus* (L.), fry, 67.
Rogan E. & Cross T. F.: Nutrient dynamics and plankton cycles in artificial ponds used in the production of oyster, *Ostrea edulis* L., spat, 9.
Rogdakis I., *see* Petridis D., 413.
Ross L. G., *see* Karplus I., 447; *see also* Martinez-Palacios C. A., 215; *see also* Martinez-Palacios C. A., 455.
Rothbard S., Rubinshtein I. & Gelman E.: Storage of common carp, *Cyprinus carpio* L., eggs for short durations, 175.
Rout S. K., *see* Mukhopadhyay P. K., 623.
Rowlerson A., *see* Nathanailides C., 885.
Rubinshtein I., *see* Rothbard S., 175.
Russell P., *see* Ojaveer H., 603.
Rye M. & Gjerde B.: Phenotypic and genetic parameters of body composition traits and flesh colour in Atlantic salmon, *Salmo salar* L., 121.

Saad C. R., *see* Das N. N., 543.
Sahu S. N. & Jana B. B.: Manipulation of stocking ratios between surface- and bottom-grazing fishes as a strategy to increase the fertilizer value of rockphosphate in a carp polyculture system, 931.
Sahul Hameed A. S., Rao P. V., Farmer J. J., Hickman-Brenner F. W. & Fanning G. R.: Characteristics and pathogenicity of a *Vibrio campbellii*-like bacterium affecting hatchery-reared *Penaeus indicus* (Milne Edwards, 1837) larvae, 853.
Salama M. E.: Effects of sex ratio and feed quality on mass production of Nile tilapia, *Oreochromis niloticus* (L.), fry, 581.
Sanchez Licea V. H., *see* Martinez-Palacios C. A., 215.
Sang Y. C., *see* Ling B. H., 775.
Sedano J., Zorrilla I., Morinigo M. A., Balebona M. C., Vidaurreta A., Bordas M. A. & Borrego J. J.: Microbial origin of the abdominal swelling affecting farmed larvae of gilthead sea bream, *Sparus aurata* L., 323.
Selosse P. M., *see* Battaglene S. C., 191.
Siddiqui A. Q., Al Hinty H. M. & Ali S. A.: Evaluation of the production potential of *Macrobranchium rosenbergii* (de Man) in monoculture and in polyculture with Nile tilapia and common carp in Saudi Arabia, 515.
Siikavuopio S. I., Baardvik B. M. & Jobling M.: Domestication effects on fin nipping, survival and

© 1996 Blackwell Science Ltd.

growth in hatchery-reared Arctic charr, *Salvelinus alpinus* (L.), in competition with wild conspecifics, 205.
Silva E. A., *see* Chellappa S., 945.
Sivakami R., Premkishore G. & Chandran M. R.: Occurrence and distribution of potentially pathogenic *Enterobacteriaceae* in carps and pond water in Tamil Nadu, India, 375.
Sivaloganathan B., Walford J., Loy G. L. & Lam T. J.: Occurrence of urinary calculi in the early larvae of brown-marbled grouper, *Epinephelus fuscoguttatus* F., 635.
Skrzypczak A., *see* Kucharczyk D., 847.
Smith A. P., *see* Young J. A., 283.
Span J. A., *see* Kamstra A., 903.
Spencer B. E., Kaiser M. J. & Edwards D. B.: The effect of Manila clam cutivation on an intertidal benthic community: the early cultivation phase, 261.
Spenser B. E., *see* Nugues M. M., 913.
Stamou E., *see* Papoutsoglou S., 25.
Steinarsson A. & Moksness E.: Oxygen consumption and ammonia excretion of common wolffish, *Anarhichas lupus* L., 1758, in an experimental-scale seawater, land-based culture system, 925.
Strand Ø.: Enhancement of bivalve production capacity in a landlocked heliothermic marine basin, 355.
Strickland N. C., *see* Nathanailides C., 885.
Strutton P. G., Bye J. A. T. & Mitchell J. G.: Determining coastal inlet flushing times: a practical expression for use in aquaculture and pollution management, 497.
Sukkel M., *see* Hariati A. M., 277.
Sureshwaran S., English W. R. & Walters E.: The profitability of producing channel catfish, *Ictalurus punctatus* (Rafinesque), in cages at a commercial scale on small farms, 557.
Svirsky F., *see* Milstein A., 79.

Tanck M. W. T., *see* Hariati A. M., 255.
Taniguchi N., *see* del Valle G., 787.
Thorarinsdóttir G. G.: Gonadal development larval settlement and growth of *Mytilus edulis* L. in a suspended population in Hvalfjördur, south-west Iceland, 57.
Toro J. E.: Phenotypic response of a cohort of *Ostrea chilensis* (Philippi 1845) to differing environmental conditions in three oyster farms in southern Chile, 1.
Treasurer J. W.: Retention of visible implant tags in farmed Atlantic salmon, *Salmo salar* L., 293.
Tsujimura A., *see* del Valle G., 787.
Tyler D. D., *see* Nathanailides C., 885.
Tziha G., *see* Papoutsoglou S., 25.

Ugedal O., *see* Heggberget T. G., 313.
Uglow R. F., *see* Buttle L. G., 391.
Utting S. D., Millican P. F. & Laing I.: The breeding potential and biochemical composition of triploid Manila clams, *Tapes philippinarum* Adams & Reeve, 573.

Vallée F., *see* Dumas S., 245.
van Dam A. A., Huisman E. A. & Rabbinge R.: Simulation of food and oxygen limitations on the growth of Nile tilapia, *Oreochromis niloticus* L., in fishponds, 463.
van der Meer M. B., Huisman E. A. & Verdegem M. C. J.: Feeding consumption, growth and protein utilization of *Colossoma macropomum* (Cuvier) at different dietary fish meal/soya meal ratios, 531.
van der Meer M. B. & Verdegem M. C. J.: Comparison of amino acid profiles of feeds and fish as a quick method for selection of feed ingredients: a case study for *Colossoma macropomum* (Cuvier), 487.
van Weerd J. H., *see* Kamstra A., 903.
Vazquez J. M., *see* Nathanailides C., 885.
Verdegem M. C. J., *see* Hariati A. M., 255; *see also* Hariati A. M., 277; *see also* van der Meer M. B., 487; *see also* van der Meer M. B., 531.
Vidaurreta A., *see* Sedano J., 323.
Vilwock W., *see* Oberst O., 235.
Vøllestad L. A., *see* Kristjánsson L. Th., 335.

Walford J., *see* Sivaloganathan B., 635.
Walters E., *see* Sureshwaran S., 557.
Wangila B. C. C. & Dick T. A.: Genetic effects and growth performance in pure and hybrid strains of rainbow trout, *Oncorhynchus mykiss* (Walbaum) (Order: Salmoniformes, Family: Salmonidae), 35
Weismann T., *see* Lahnsteiner F., 659.
Weyl O., Kaiser H. & Hecht T.: On the efficacy and mode of action of 2-phenoxyethanol as an anaesthetic for goldfish, *Carassius auratus* (L.), at different temperatures and concentrations, 757.
Wheatley S. B., *see* Menzies F. D., 183.
Wiadnya D. G. R., *see* Hariati A. M., 255; *see also* Hariati A. M., 277.
Woodward K. N.: The regulation of fish medicines—UK and European Union aspects, 725.
Wyszomirska E., *see* Kucharczyk D., 847.

Young J. A., Smith A. P. & Muir J. F.: Market analysis using Multiple Correspondence Analysis: the case of consumer acceptance of African catfish, 283.

Zakes Z. & Demska-Zakes K.: Effect of diets on growth and reproductive development of juvenile pikeperch, *Stizostedion lucioperca* (L.), reared under intensive culture conditions, 841.
Zhou L. H., *see* Guan R. Z., 587.
Zorrilla I., *see* Sedano J., 323.

Subject Index

Abdominal swelling, *Vibrio* spp., gilthead sea bream, 323.
Acipenser fulvescens Rafinesque, *see* lake sturgeon.
Adaptability, haemoglobin, tilapia, 597.
Adhesiveness, egg swelling, wolffish, 421.
Algae, light administration, Atlantic halibut, 101.
Algal concentration, filtration and ingestion rate, carpet shell clam, 613.
Alumino-silicate clays, zeolites, water quality, 301.
Amino acid profiles:
  Colossoma macropomum, diet, 487;
  diet:
    Colossoma macropomum, 487;
    prawn, 543;
  prawn, diet, 543.
Ammonia excretion:
  Atlantic salmon, diet, 937;
  diet, Atlantic salmon, 937;
  oxygen consumption, wolffish, 925;
  wolffish, oxygen consumption, 925.
Anaesthetic, 2-phenoxyethanol, goldfish, 757.
Anarhichas lupus L., *see* wolffish.
Anguilla anguilla L., *see* eel.
Animal by-product meal, diet, Nile tilapia, 67.
Argopecten circularis (Sowerby), *see* Catarina scallop.
Artificial propagation, fertilization, eel, 587.
Ascorbate status, gonad development, yellow perch, 539.
Ayu, genetic differences, haematological traits, 787.

Bass:
  Australian:
    hormone inducement, ovulation and spawning, 191;
    ovulation and spawning, hormone inducement, 191;
  sea:
    ectoparasites, gilthead sea bream, environmental conditions, 25;
    environmental conditions, ectoparasites, gilthead sea bream, 25;
    gilthead sea bream, environmental conditions, ectoparasites, 25;
    muscle cellularity, somatic growth, 885;
    somatic growth, muscle cellularity, 885.
Batch mark, elastomer, injection, 631.
Benthic community changes, intertidal cultivation, Pacific oyster, 913.
Biochemical and immunological markers, discrimination, tilapia, 235.
Biochemical composition, breeding potential, Manila clam, 573.
Biomass, plankton production, fertilizer, 157.
Bivalve production, heliothermic marine basin, 355.
Body composition traits, flesh colour, Atlantic salmon, 121.
Branchionus calyciflorus Pallas, *see* rotifer.
Bream, gilthead sea:
  abdominal swelling, *Vibrio* spp., 323;
  diet, soybean meal, 147;
  ectoparasites, environmental conditions, sea bass, 25;
  environmental conditions, ectoparasites, sea bass, 25;
  growth models, 413;
  growth rate, salinity, 763;
  sea bass, environmental conditions, ectoparasites, 25.
  salinity, growth rate, 763.
  soybean meal, diet, 147.
  stress, vitamin C status, 405;
  Vibrio spp., abdominal swelling, 323;
  vitamin C status, stress, 405.
Breeding potential, biochemical composition, Manila clam, 573.

Cage culture, profitability, channel catfish, 557.
Carassius auratus (L.), *see* goldfish.
^{14}C-glucose injection, dietary carbohydrate, Atlantic halibut, 565.
Carp pituitary extract, human chorionic gonadotrophin, induced spawning, perch, 847.
Carp:
  common:
    diet, growth response, 673;
    eggs, short-term storage, 175;
    Enterobacteriaceae, water quality, 375;
    growth response, diet, 673;
    production characteristics and profitability, stocking ratio, 447;
    rockphosphate, stocking ratio, 931;
    short-term storage, eggs, 175;
    stocking ratio:
      rockphosphate, 931;
      production characteristics and profitability, 447;
    water quality, Enterobacteriaceae, 375;
  Indian major:
    diet, lipid supplements, 623;
    growth curve analysis, Hill's procedure, 891;
    Hill's procedure, growth curve analysis, 891;
    lipid supplements, diet, 623.
Catfish:
  African:
    diet:
      nitrogen efflux rates, temperature, 391;
      oxidative nutritional stress, 795;
      temperature, nitrogen efflux rates, 391;
    market analysis, Multiple Correspondence Analysis, 283;
    Multiple Correspondence Analysis, market analysis, 283;
    nitrogen efflux rates, temperature, diet, 391;
    oxidative nutritional stress, diet, 795;
    temperature, diet, nitrogen efflux rates, 391;
  channel:

© 1996 Blackwell Science Ltd.

cage culture, profitability, 557;
profitability, cage culture, 557;
estuarine:
feeding frequency, water quality, 107;
water quailty, feeding frequency, 107;
tropical:
diet, growth, 823;
growth, diet, 823.
Catla catla (Hamilton), *see* Indian major carp.
Charr:
Arctic:
Atlantic salmon, diet, Na+L-lactate, 429;
brook charr, hybrids, survival, 245;
demand feeding behaviour, diet, 479;
diet:
Atlantic salmon, Na+L-lactate, 429;
demand feeding behaviour, 479;
Na^+-lactate, Atlantic salmon, 429;
domestication effects, fin nipping, 205;
fin nipping, domestication effects, 205;
hybrids, brook charr, survival, 245;
Na^+-lactate, diet, Atlantic salmon, 429;
sea ranching, stress, 735;
stress, sea ranching, 735;
survival, hybrids, brook charr, 245;
brook:
Arctic charr, hybrids survival, 245;
hybrids, Arctic charr, survival, 245;
survival, hybrids, Arctic charr, 245.
Chelon labrosus (Risso), *see* thick-lipped grey mullet.
Chrysichthys nigrodigitatus Lacépède, *see* estuarine catfish.
Cichlasoma synspilum:
diet, protein requirements, 167;
protein requirements, diet, 167;
respiratory characteristics, salinity tolerance, 215;
salinity tolerance, respiratory characteristics, 215.
Cichlasoma urophthalmus, temperature, food intake, 455.
Clam:
carpet shell:
algal concentration, filtration and ingestion rate, 613;
culture methods, growth, 399;
filtration and ingestion rate, algal concentration, 613;
growth, culture methods, 399;
Manila:
biochemical composition, breeding potential, 573;
breeding potential, biochemical composition, 573;
early cultivation phase, intertidal benthic community, 261;
intertidal benthic community, early cultivation phase, 261.
Clarias gariepinus (Burchell), *see* African catfish.
Clariid catfish, African, testes, sperm, cryopreservation, 805.
Coastal inlets, flushing, model, 497.
Colossoma macropomum (Cuvier), *see* tambaqui.
Competition, diet, hybrid red tilapia, 945.
Computerized information retrieval system, disease control, Atlantic salmon, 183.
CPE and HCG treatments, induced breeding, tambaqui, 91.
Crassostrea gigas Thunberg, *see* Pacific oyster.
Cryopreservation, testes, sperm, African clariid catfish, 805.
Culture methods:
carpet shell clams, growth, 399;
growth, carpet shell clams, 399;
prawn, production potential, 515;
production potential, prawn, 515.
Cyprinus carpio L., *see* carp.
Cytochalasin B and 6-dimethylaminopurine, triploidy, Sydney rock oyster, 689.
Cytological blood parameters, leucogram, rainbow trout, 437.

Demand feeding behaviour, diet, Arctic charr, 479.
Dicentrarchus labrax (L.), *see* sea bass.
Diet:
African catfish:
nitrogen efflux rates, temperature, 391;
oxidative nutritional stress, 795;
temperature, nitrogen efflux rates, 391;
amino acid profiles:
prawn, 543;
tambaqui, 487;
ammonia excretion, Atlantic salmon, 937;
animal by-product meal, Nile tilapia, 67;
Arctic charr:
Atlantic salmon, Na^+-lactate, 429;
demand feeding behaviour, 479;
Na^+-lactate Atlantic salmon, 429;
Atlantic salmon:
ammonia excretion, 937;
Arctic charr, Na^+-lactate, 429;
Na^+-lactate, Arctic charr, 429;
Cichlasoma synspilum, protein requirements, 167;
common carp, growth response, 673;
competition, hybrid red tilapia, 945;
demand feeding behaviour, Arctic charr, 479;
gilthead sea bream, soybean meal, 147;
Gracilaria sp. meal, shrimp, 345;
growth:
pikeperch, 841;
tropical catfish, 823;
growth performance, Nile tilapia, 523.
growth rate, tambaqui, 531.
growth response, common carp, 673.
hybrid red tilapia, competition, 945;
Indian major carp, lipid supplements, 623;
lake sturgeon, stimulants, 953;
lipid supplements, Indian major carp, 623;
Na^+-lactate, Arctic charr, Atlantic salmon, 429;
Nile tilapia:
animal by-product meal, 67;
growth performance, 523;
nitrogen efflux rates, temperature, African catfish, 391;
oxidative nutritional stress, African catfish, 795;
oxygen limitations, Nile tilapia, 463;
pikeperch, growth, 841;
prawn, amino acid profiles, 543;
protein requirements, *Cichlasoma synspilum*, 167;
protein-to-energy ratio, thick-lipped grey mullet, 603;

© 1996 Blackwell Science Ltd.

rainbow trout, water quality, 699;
shrimp, *Gracilaria* sp. meal, 345;
soybean meal, gilthead sea bream, 147;
stimulants, lake sturgeon, 953;
tambaqui:
amino acid profiles, 487;
tambaqui, 531;
temperature, nitrogen efflux rates, African catfish, 391;
thick-lipped grey mullet, protein-to-energy ratio, 603;
tropical catfish, growth, 823;
water quality, rainbow trout, 699.
Dietary carbohydrate, ^{14}C-glucose injection, Atlantic halibut, 565.
Discrimination, biochemical and immunological markers, tilapia, 235.
Disease control, computerized information retrieval system, Atlantic salmon, 183.
Domestication effects, fin nipping, Arctic charr, 205.

Early cultivation phase, intertidal benthic community, Manila clam, 261.
Ectoparasites, environmental conditions, European sea bass, gilt-head sea-bream, 25.
Eel:
artificial propagation, fertilization, 587;
feeding stimulation, rapid-screening method, 379;
fertilization, artificial propagation, 587;
nitrite, toxicity, 903;
rapid-screening method, feeding stimulation, 379;
toxicity, nitrite, 903.
Effluent:
extracellular-enzyme activity, trout, 895;
mangrove seedling shoot biomass, shrimp, 745;
shrimp, mangrove seedling shoot biomass, 745;
trout, extracellular-enzyme activity, 895.
Egg swelling, adhesiveness, wolffish, 421.
Eggs, short-term storage, common carp, 175.
Elastomer, injection, batch mark, 631.
Energy budget, ontogeny, sole, 831.
Enterobacteriaceae, common carp, water quality, 375.
Environmental conditions:
Chilean oyster phenotypic response, 1;
ectoparasites, European sea bass, gilthead sea bream, 25;
gilthead sea bream, sea bass, ectoparasites, 25;
phenotypic response, Chilean oyster, 1;
sea bass, gilthead sea bream, ectoparasites, 25.
Epinephelus fuscoguttatus F., *see* brown-marbled grouper.
Experimental culture conditions, nitrogen budget, white steenbras, 43.
Export performance, Japan, USA, shrimp, 775.
Extracellular-enzyme activity, effluent, trout, 895.

Fatty acid profiles, rotifer, gudgeon, perch, 651.
Feed quality, sex ratio, Nile tilapia, 581.
Feeding frequency, water quality, catfish, 107.
Feeding stimulation, rapid-screening method, eel, 379.
Fertilization:
artificial propagation, eel, 587;
eel, artificial propagation, 587;
incubation, scallop, 505;
salmonids, semen crypreservation, 659;
semen cryopreservation, salmonids, 659;
scallop, incubation, 505.
Fertilizer, plankton production, biomass, 157.
Fin nipping, domestication effects, Arctic charr, 205.
Flesh colour, body composition traits, Atlantic salmon, 121.
Filtration and ingestion rate, algal concentration, carpet shell clam, 613.
Flushing, coastal inlets, model, 497.
Food intake, temperature, *Cichlasoma urophthalmus*, 455.

Genetic differences:
ayu, haematological traits, 787;
haematological traits, ayu, 787;
rainbow trout, starch gel electrophoresis, 643;
starch gel electrophoresis, rainbow trout, 643.
Genetic effects, growth performance, rainbow trout, 35.
Gobio gobio (L.), *see* gudgeon.
Goldfish, 2-phenoxyethanol, anaesthetic, 757.
Gonad development:
ascorbate status, yellow perch, 539;
larval settlement, mussel, 57;
mussel, larval settlement, 57;
yellow perch, ascorbate status, 539.
Gracilaria sp. meal, diet, shrimp, 345.
Grouper, brown-marbled, urinary calculi, 635.
Growth:
carpet shell clams, culture methods, 399;
Caterina scallop, stocking density, 711;
Colossoma macropomum, diet, 531;
common carp, diet, 673;
culture methods, carpet shell clams, 399;
diet:
Colossoma macropomum, 531;
common carp, 673;
pikeperch, 841;
Nile tilapia, 523;
tropical catfish, 823;
genetic effects, rainbow trout, 35;
gilthead sea bream, salinity, 763;
Nile tilapia, diet, 523;
pikeperch, diet, 841;
rainbow trout, genetic effects, 35;
salinity, gilthead sea bream, 763;
stocking density, Caterina scallop, 711.
tropical catfish, diet, 823.
Growth curve analysis, Hill's procedure, Indian major carp, 891.
Growth dynamics, sexual maturation, brown trout, 815.
Growth models, sea bream, 413.
Gudgeon, perch, rotifer, fatty acid profiles, 651.

Haematological traits, genetic differences, ayu, 787.
Haemoglobin, adaptability, tilapia, 597.
Halibut, Atlantic:
algae, light administration, 101;
^{14}C-glucose injection, dietary carbohydrate, 565;
dietary carbohydrate, ^{14}C-glucose injection, 565;
light administration, algae, 101.
Heliothermic marine basin, bivalve production, 355.
Hepatic enzymes, selenium dietary enhancement, coho salmon, 135.

Heterobranchus longifilis (Valenciennes), *see* African clariid catfish.
Hill's procedure, growth curve analysis, Indian major carp, 891.
Hippoglossus hippoglossus (L.), *see* Atlantic halibut.
Hormone inducement, ovulation and spawning, Australian bass, 191.
Human chorionic gonadotrophin, carp pituitary extract, induced spawning, perch, 847.
Hybrids, survival, brook and Arctic charr, 245.

Ictalurus punctatus (Rafinesque), *see* channel catfish.
Incubation, fertilization, scallop, 505.
Indigenous species, polyculture, Malawi, 225.
Individual variation, progeny size and quality, rainbow trout, 335.
Indonesia:
  pond production, shrimp, 277;
  shrimp:
    pond production, 277;
    water quality, 255;
  water quality, shrimp, 255.
Induced breeding, CPE and HCG treatments, tambaqui, 91.
Induced spawning, carp pituitary extract, human chorionic gonadotrophin, perch, 847.
Ingestion efficiency, live feed, giant prawn, 113.
Injection, elastomer, batch mark, 631.
Intertidal benthic community, early cultivation phase, Manila clam, 261.
Intertidal cultivation, benthic community changes, Pacific oyster, 913.

Japan, USA, export performance, shrimp, 775.

Larval settlement, gonad development, mussel, 57.
Leucogram, cytological blood parameters, rainbow trout, 437.
Life-cycle, management, wolffish, 865.
Light administration, algae, Atlantic halibut, 101.
Lipid supplements, diet, Indian major carp, 623.
Lithognathus lithognathus Cuvier, *see* white steenbras.
Live feed, ingestion efficiency, giant prawn, 113.
Lutenizing hormone-releasing hormone analogue, sperm count Nile tilapia, 95.

Macquaria novemaculeata (Steindachner), *see* Australian bass.
Macrobranchium rosenbergii (de Man), *see* giant prawn.
Malawi, polyculture, indigenous species, 225.
Management, life-cycle, wolffish, 865.
Mangrove seedling shoot biomass, effluent, shrimp, 745.
Market analysis, Multiple Correspondence Analysis, African catfish, 283.
Model, flushing, coastal inlets, 497.
Monopterus albus (Zuiew), *see* eel.
Mullet, thick-lipped grey, protein-to-energy ratio, diet, 603.
Multiple Correspondence Analysis, market analysis, African catfish, 283.
Muscle cellularity, somatic growth, sea bass, 885.
Mussel, gonad development, larval settlement, 57.
Mystus nemurus (Cuvier & Valenciennes), *see* tropical catfish.
Mytilus edulis L., *see* mussel.

Na^+-lactate, diet, Arctic charr, Atlantic salmon, 429.
Nitrite, toxicity, eel, 903.
Nitrogen budget, experimental culture conditions, white steenbras, 43.
Nitrogen efflux rates, temperature, diet, African catfish, 391.
Nutrient dynamics, plankton cycles, European flat oyster, 9.

Oncorhynchus kisutch (Walbaum), *see* coho salmon.
Oncorhynchus mykiss (Walbaum), *see* rainbow trout.
Ontogeny, energy budget, sole, 831.
Oreochromis niloticus (L.), *see* Nile tilapia.
Oreochromis niloticus (L.) × *O. mossambicus* (Peters), *see* hybrid red tilapia.
Ostrea chilensis (Philippi 1845), *see* Chilean oyster.
Ostrea edulis L., *see* European flat oyster.
Ovulation and spawning, hormone inducement, Australian bass, 191.
Oxidative nutritional stress, diet, African catfish, 795.
Oxygen consumption, ammonia excretion, wolffish, 925.
Oxygen limitations, diet, Nile tilapia, 463.
Oyster:
  Chilean:
    environmental conditions, phenotypic response, 1;
    phenotypic response, environmental conditions, 1;
  European flat:
    nutrient dynamics, plankton cycles, 9;
    plankton cycles, nutrient dynamics, 9;
  Pacific:
    benthic community changes, intertidal cultivation, 913;
    intertidal cultivation, benthic community changes, 913;
  Sydney rock:
    cytochalasin B and 6-dimethylaminopurine, triploidy, 689;
    triploidy, cytochalasin B and 6-dimethylaminopurine, 689.

Pathogenicity, *Vibrio campbellii*-like bacterium, shrimp, 853.
Pecten fumatus Reeve, *see* scallop.
Penaeus indicus (Milne Edwards), *see* shrimp.
Penaeus monodon Fabricius, *see* shrimp.
Perca flavescens (Mitchill), *see* yellow perch.
Perca fluviatilis L., *see* perch.
Perch:
  carp pituitary extract, human chorionic gonadotrophin, induced spawning, 847;
  fatty acid profiles, gudgeon, rotifer, 651;
  gudgeon, rotifer, fatty acid profiles, 651;
  human chorionic gonadotrophin, carp pituitary extract, induced spawning, 847;
  induced spawning, carp pituitary extract, human chorionic gonadotrophin, 847;
  rotifer, gudgeon, fatty acid profiles, 651;
  yellow:

© 1996 Blackwell Science Ltd, *Aquaculture Research*, **27**, 965–970

ascorbate status, gonad development, 539;
gonad development, ascorbate status, 539.
Phenotypic response, Chilean oyster, environmental conditions, 1.
2-Phenoxyethanol, anaesthetic, goldfish, 757.
Pikeperch, growth, diet, 841.
Plankton:
composition:
species combinations, water chemistry, 79;
water chemistry, species combinations, 79;
cycle:
European flat oyster, nutrient dynamics, 9;
nutrient dynamics, European flat oyster, 9;
production:
biomass, fertilizer, 157;
fertilizer, biomass, 157.
Plecoglossus altivelis Temminck & Schlegel, *see* ayu.
Polyculture, indigenous species, Malawi, 225.
Pond depth, water temperature, Nile tilapia, 681.
Pond production, shrimp, Indonesia, 277.
Prawn:
amino acid profiles, diet, 543;
culture methods, production potential, 515;
diet, amino acid profiles, 543;
production potential, culture methods, 515;
giant:
ingestion efficiency, live feed, 113;
live feed, ingestion efficiency, 113.
Prespawning migratory behaviour, Atlantic salmon, 313.
Production characteristics and profitability, stocking ratio, common carp, 447.
Production potential, culture methods, prawn, 515.
Profitability, cage culture, channel catfish, 557.
Progeny size and quality, individual variation, rainbow trout, 335.
Protein requirements, diet, *Cichlasoma synspilum*, 167.
Protein-to-energy ratio, diet, thick-lipped grey mullet, 603.

Rapid-screening method, feeding stimulation, eel, 379.
Regulation, veterinary medicine, UK and European Union, 725.
Respiratory characteristics, salinity tolerance, *Cichlasoma synspilum*, 215.
Rockphosphate, stocking ratio, common carp, 931.
Rotifer, fatty acid profiles, gudgeon, perch, 651.

Saccostrea commercialis (Iredale & Roughley), *see* Sydney rock oyster.
Salinity tolerance, respiratory characteristics, *Cichlasoma synspilum*, 215.
Salinity, growth rate, gilthead sea bream, 763.
Salmo salar L., *see* Atlantic salmon.
Salmon:
Atlantic:
ammonia excretion, diet, 937;
Arctic charr, diet, Na+1-lactate, 429;
body composition traits, flesh colour, 121;
computerized information retrieval system, disease control, 183;
diet:
ammonia excretion, 937;
Arctic charr, Na^+-lactate, 429;
Na^+-lactate, Arctic charr, 429;
disease control, computerized information retrieval system, 183;
flesh colour, body composition traits, 121;
Na^+-lactate, Arctic charr, diet, 429;
prespawning migratory behaviour, 313;
visible implant tags, 293;
coho:
hepatic enzymes, selenium dietary enhancement, 135;
selenium dietary enhancement, hepatic enzymes, 135.
Salmonids, semen cryopreservation, fertilization rate, 659.
Salvelinus alpinus (L.), *see* Arctic charr.
Salvelinus fontinalis (Mitchill), *see* brook charr.
Scallop:
Caterina:
growth rate, stocking density, 711;
stocking density, growth rate, 711;
fertilization, incubation, 505;
incubation, fertilization, 505.
Sea ranching, stress, Arctic charr, 735.
Selenium dietary enhancement, hepatic enzymes, coho salmon, 135.
Semen cryopreservation, fertilization rate, salmonids, 659.
Sex ratio, feed quality, Nile tilapia, 581.
Sexual maturation, growth dynamics, brown trout, 815.
Short-term storage, eggs, common carp, 175.
Shrimp:
diet, *Gracilaria* sp. meal, 345;
effluent, mangrove seedling shoot biomass, 745;
export performance, Japan, USA, 775;
Gracilaria sp. meal, diet, 345;
Indonesia:
pond production, 277;
water quality, 255;
Japan, USA, export performance, 775;
mangrove seedling shoot biomass, effluent, 745;
pathogenicity, *Vibrio campbellii*-like bacterium, 853;
pond production, Indonesia, 277;
USA, Japan, export performance, 775;
Vibrio campbellii-like bacterium, pathogenicity, 853;
water quality, Indonesia, 255;
Sole, ontogeny, energy budget, 831.
Solea solea (L.), *see* sole.
Somatic growth, muscle cellularity, sea bass, 885.
Soybean meal, diet, gilthead sea bream, 147.
Sparus aurata L., *see* gilthead sea bream.
Species combinations, water chemistry, plankton composition, 79.
Sperm, testes, cryopreservation, African clariid catfish, 805.
Sperm count, lutenizing hormone-releasing hormone analogue, Nile tilapia, 95.
Starch gel electrophoresis, genetic differences, rainbow trout, 643.
Steenbras, white, nitrogen budget, experimental culture conditions, 43.

Stimulants, diet, lake sturgeon, 953.
Stizostedion lucioperca (L.), *see* pikeperch.
Stocking density, growth rate, Caterina scallop, 711.
Stocking ratio:
  common carp:
    rockphosphate, 931;
    production characteristics and profitability, 447;
  production characteristics and profitability, common carp, 447;
  rockphosphate, common carp, 931;
Stress:
  Arctic charr, sea ranching, 735;
  gilthead sea bream, vitamin C status, 405;
  sea ranching, Arctic charr, 735;
  vitamin C status, gilthead sea bream, 405.
Sturgeon, lake, diet, stimulants, 953.
Survival, hybrids, brook and Arctic charr, 245.

Tambaqui:
  amino acid profiles, diet, 487;
  CPE and HCG treatments, induced breeding, 91;
  diet:
    amino acid profiles, 487;
    growth rate, 531;
  growth rate, diet, 531;
  induced breeding, CPE and HCG treatments, 91.
Tapes decussatus (L.), *see* carpet shell clam.
Tapes philippinarum Adams & Reeve, *see* Manila clam.
Temperature:
  African catfish, nitrogen efflux rates, diet, 391;
  Cichlasoma urophthalmus, food intake, 455;
  diet, nitrogen efflux rates, African catfish, 391;
  food intake, *Cichlasoma urophthalmus*, 455;
  nitrogen efflux rates, diet, African catfish, 391.
Testes, sperm, cryopreservation, African clariid catfish, 805.
Tilapia:
  adaptability, haemoglobin, 597;
  biochemical and immunological markers, discrimination, 235;
  discrimination, biochemical and immunological markers, 235;
  haemoglobin, adaptability, 597;
  hybrid red:
    competition, diet, 945;
    diet, competition, 945;
  Nile:
    animal by-product meal, diet, 67;
    diet:
      animal by-product meal, 67;
      growth performance, 523;
      oxygen limitations, 463;
    feed quality, sex ratio, 581;
    growth performance, diet, 523;
    lutenizing hormone-releasing hormone analogue, sperm count, 95;
    oxygen limitations, diet, 463;
    pond depth, water temperature, 681;
    sex ratio, feed quality, 581;
    sperm count, lutenizing hormone-releasing hormone analogue, 95;
    water temperature, pond depth, 681.
Toxicity, nitrite, eel, 903.
Triploidy, cytochalasin B and 6-dimethylaminopurine, Sydney rock oyster, 689.
Trout:
  brown:
    growth dynamics, sexual maturation, 815;
    sexual maturation, growth dynamics, 815;
  effluent, extracellular-enzyme activity, 895;
  extracellular-enzyme activity, effluent, 895;
  rainbow:
    cytological blood parameters, leucogram, 437;
    diet, water quality, 699;
    genetic differences, starch gel electrophoresis, 643;
    genetic effects, growth performance, 35;
    growth performance, genetic effects, 35;
    individual variation, progeny size and quality, 335;
    leucogram, cytological blood parameters, 437;
    progeny size and quality, individual variation, 335;
    starch gel electrophoresis, genetic differences, 643;
    water quality, diet, 699.

UK and European Union, regulation, veterinary medicine, 725.
Urinary calculi, brown-marbled grouper, 635.
USA, Japan, export performance, shrimp, 775.

Veterinary medicine, regulation, UK and European Union, 725.
Vibrio spp., abdominal swelling, gilthead sea bream, 323.
Vibrio campbellii-like bacterium, pathogenicity, shrimp, 853.
Visible implant tags, Atlantic salmon, 293.
Vitamin C status, stress, gilthead sea bream, 405.

Water chemistry, plankton composition, species combinations, 79.
Water quality:
  alumino-silicate clays, zeolites, 301;
  catfish, feeding frequency, 107;
  common carp, Enterobacteriaceae, 375;
  diet, rainbow trout, 699;
  Enterobacteriaceae, common carp, 375;
  feeding frequency, catfish, 107;
  Indonesia, shrimp, 255;
  shrimp, Indonesia, 255;
  rainbow trout, diet, 699;
  zeolites, alumino-silicate clays, 301.
Water temperature, pond depth, Nile tilapia, 681.
Wolffish:
  adhesiveness, egg swelling, 421;
  ammonia excretion, oxygen consumption, 925;
  egg swelling, adhesiveness, 421;
  life-cycle, management, 865;
  management, life-cycle, 865;
  oxygen consumption, ammonia excretion, 925.

Zeolites, alumino-silicate clays, water quality, 301.

© 1996 Blackwell Science Ltd.